

What is AVMAPKI FAKZYNJA CO-PACK?

AVMAPKI FAKZYNJA CO-PACK is a prescription medicine used to treat adults who have low-grade serous ovarian cancer (LGSOC) that has come back (recurrent), **and** LGSOC with an abnormal *KRAS* gene, **and** previously been treated with medicine for their cancer.

It is not known if AVMAPKI FAKZYNJA CO-PACK is safe and effective in children.



NOW APPROVED

AVMAPKI FAKZYNJA CO-PACK

The first and only prescription medicine used to treat adults who have low-grade serous ovarian cancer (LGSOC) that has come back (recurrent), and LGSOC with an abnormal *KRAS* gene, who have previously been treated with medicine for their cancer.

Click to read full [Prescribing Information](#) and [Patient Information](#).

WHAT IS AVMAPKI FAKZYNJA CO-PACK?

AVMAPKI FAKZYNJA CO-PACK is a prescription medicine used to treat adults who have low-grade serous ovarian cancer (LGSOC) that has come back (recurrent), and LGSOC with an abnormal *KRAS* gene, who have previously been treated with medicine for their cancer. It is not known if AVMAPKI FAKZYNJA CO-PACK is safe and effective in children.

WHAT SHOULD I KNOW BEFORE TAKING AVMAPKI FAKZYNJA CO-PACK?

Before taking AVMAPKI FAKZYNJA CO-PACK, be sure to tell your healthcare provider about all of your medical conditions, including if you:

- have a history of eye or vision problems
- have had severe skin reactions
- have liver problems
- are pregnant or plan to become pregnant, as AVMAPKI FAKZYNJA CO-PACK can harm your unborn baby


Females who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment
- You should use effective birth control (contraception) during treatment and for 1 month after the last dose
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment

Males with a female partner who is able to become pregnant:

- You should use effective birth control (contraception) during treatment and for 4 months after the last dose
- are breastfeeding or plan to breastfeed. It is not known if AVMAPKI FAKZYNJA CO-PACK passes into your breast milk. Do not breastfeed during treatment and for 2 weeks after the last dose

Be sure to tell your healthcare provider about all the prescription and over-the-counter medicines, vitamins, and herbal supplements you take, especially proton pump inhibitors (PPIs) or H2 receptor antagonists, and a blood thinner called warfarin.



Click to read full
[Prescribing Information](#)
and [Patient Information](#).

 **AVMAPKI™ FAKZYNJA™ CO-PACK**
(avutemetinib capsules; defactinib tablets) 0.8mg/200mg

AVMAPKI FAKZYNJA CO-PACK may affect the way other medicines work, and other medicines may affect how AVMAPKI FAKZYNJA CO-PACK works.

HOW SHOULD I TAKE AVMAPKI FAKZYNJA CO-PACK?

AVMAPKI:

- On **weeks 1, 2, and 3**, take 3.2 mg (4 capsules) on **day 1** and **day 4**



3.2 mg
4 capsules = 1 dose

- On **week 4**, take a one-week break
- Take each dose at the same time **with food**
- Swallow the capsules whole. Do not chew, break, or open the capsules
- **Keep refrigerated** in original bottle

FAKZYNJA:

















- On **weeks 1, 2, and 3**, take 200 mg (1 tablet) **2 times a day**



200 mg
1 tablet = 1 dose

- On **week 4**, take a one-week break
- Take **with food**
- Swallow the tablets whole. Do not chew, break, or crush the tablets
- **Keep refrigerated** in original bottle

Use the table below for monthly treatment guidance.

	Time	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Weeks 1, 2, & 3	Morning	 + 			 + 			
	Evening							
Week 4	One-week break. Do not take AVMAPKI FAKZYNJA CO-PACK during week 4.							

Images shown are not actual size

Take AVMAPKI FAKZYNJA CO-PACK exactly as your healthcare provider tells you to and do not change your dose or stop taking it without talking to your healthcare provider. If you develop certain side effects, your healthcare provider may change your dose, temporarily stop treatment, or completely stop treatment.

Click to read full
[Prescribing Information](#)
and [Patient Information](#).

 **AVMAPKI™ FAKZYNJA™ CO-PACK**
(avutometinib capsules; defactinib tablets) 0.8 mg/200 mg

CAN I TAKE AVMAPKI FAKZYNJA CO-PACK WITH ANTACIDS?

If you take antacids, make sure you take FAKZYNJA tablets 2 hours before or 2 hours after taking the antacid.

WHAT SHOULD I DO IF I MISS A DOSE?

If you miss a dose of AVMAPKI by more than 24 hours, skip the missed dose and take the next scheduled dose as prescribed by your healthcare provider. Do not take two doses at the same time to make up for a missed dose.

If you miss a dose of FAKZYNJA by more than 6 hours, skip the missed dose and take the next scheduled dose as prescribed by your healthcare provider. Do not take two doses at the same time to make up for a missed dose.

If you vomit after taking AVMAPKI capsules or FAKZYNJA tablets, do not take an additional dose. Take the next scheduled dose as prescribed by your healthcare provider.

IS THERE ANYTHING ELSE I SHOULD KNOW?

Before and during treatment with AVMAPKI FAKZYNJA CO-PACK, your healthcare provider will give you medicines to apply to your face, scalp, neck, upper chest, and upper back, and antibiotic medicines to take by mouth to help prevent skin reactions.

WHAT SHOULD I AVOID WHILE TAKING AVMAPKI FAKZYNJA CO-PACK?

Limit your time in the sun. AVMAPKI FAKZYNJA CO-PACK can make your skin sensitive to the sun and the light from sunlamps and tanning beds, which means you may burn more easily and get a severe sunburn, so be sure to apply daily sunscreen with a Sun Protection Factor (SPF) of 30 or more. Wear a hat and clothes that cover your skin to protect against sunburn. Tell your healthcare provider if you get a sunburn.



WHAT ARE THE POSSIBLE SIDE EFFECTS OF AVMAPKI FAKZYNJA CO-PACK?

AVMAPKI FAKZYNJA CO-PACK may cause serious side effects, including:

- **Eye problems.** Eye problems are common with AVMAPKI FAKZYNJA CO-PACK and can be severe. Your healthcare provider will send you to see an eye care professional before starting treatment, during treatment, and for any new or worsening eye problems. Tell your healthcare provider if you get any new or worsening eye problems, including:
 - changes in vision, such as blurred vision, double vision, or vision loss
 - dry eye(s)
 - eye and eyelid inflammation, including redness, swelling, and itching
 - eye pain
 - new or increased floaters (small dark spots or squiggly lines that float across your vision)
 - light hurting eyes or seeing flashes of light

AVMAPKI FAKZYNJA CO-PACK may cause serious side effects, including:

- **Severe skin reactions.** Skin reactions are common with AVMAPKI FAKZYNJA CO-PACK and can be severe or life threatening. Tell your healthcare provider if you develop any new or worsening skin reactions, including:
 - skin rash or acne
 - dry skin
 - itching
 - blisters on your skin
 - redness or swelling of your face, hands, or soles of your feet
 - blisters or sores in your mouth
 - peeling of your skin
 - skin infection
- **Liver problems.** Liver problems are common with AVMAPKI FAKZYNJA CO-PACK and can be severe. Treatment with AVMAPKI FAKZYNJA CO-PACK can increase the level of aspartate aminotransferase, alanine aminotransferase, and bilirubin in your blood. Your healthcare provider will do blood tests before starting each treatment cycle and during treatment to check your liver function. Tell your healthcare provider if you get any signs or symptoms of severe liver problems including:
 - yellowing of your skin or the white part of your eyes
 - itchy skin
 - feeling very tired
 - flu-like symptoms
 - nausea or vomiting
 - upper right side stomach pain
 - dark or tea colored urine
 - bleeding, or bruising more easily than normal
- **Muscle problems (rhabdomyolysis).** AVMAPKI FAKZYNJA CO-PACK may cause muscle problems that can be severe. Treatment with AVMAPKI FAKZYNJA CO-PACK can increase the level of an enzyme in your blood called creatine phosphokinase (CPK) and can be a sign of muscle damage. Your healthcare provider will perform blood tests to check your levels of CPK before starting each treatment cycle and during treatment. Tell your healthcare provider if you get any signs or symptoms of increased CPK including:
 - weakness or difficulty moving arms and legs
 - muscle or bone aches or pain that does not go away
 - dark, red colored urine or decreased urine output

Click to read full
[Prescribing Information](#)
and [Patient Information](#).



AVMAPKI™ FAKZYNJA™ CO-PACK
(avutometinib capsules; defactinib tablets) 0.8mg/200mg

The most common side effects of AVMAPKI FAKZYNJA CO-PACK include:

- nausea
- tiredness
- increased aspartate aminotransferase
- diarrhea
- swelling in the body (edema)
- decreased hemoglobin
- increased alanine aminotransferase
- vomiting
- increased blood bilirubin
- increased triglycerides
- stomach-area pain
- acid reflux
- increased alkaline phosphatase
- decreased platelets
- constipation
- shortness of breath
- cough
- urinary tract infection
- decreased lymphocytes
- decreased neutrophils
- proteinuria

CAN I CONCEIVE WHILE TAKING AVMAPKI FAKZYNJA CO-PACK?

AVMAPKI FAKZYNJA CO-PACK may cause fertility problems in females and males, which may affect your ability to have children. If this is a concern for you, be sure to talk to your healthcare provider.

WHAT SHOULD I DO IF I HAVE SIDE EFFECTS?

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of AVMAPKI FAKZYNJA CO-PACK. Call your healthcare provider for medical advice about side effects.

You may report side effects to the FDA at 1-800-FDA-1088.

HOW SHOULD I STORE AVMAPKI FAKZYNJA CO-PACK?

You should keep your AVMAPKI capsules and FAKZYNJA tablets in their original, child-resistant bottles and store them in the refrigerator between 36°F and 46°F (2°C to 8°C).

Always be sure to **keep AVMAPKI FAKZYNJA CO-PACK and all medicines out of the reach of children.**

WANT TO KNOW MORE ABOUT AVMAPKI FAKZYNJA CO-PACK?

Talk to your healthcare provider and visit AvmapkiFakzynjaCo-Pack.com

GET THE SUPPORT YOU NEED TODAY

We understand healthcare can be complex, so we're here to help you access AVMAPKI FAKZYNJA CO-PACK as prescribed by your healthcare provider. Regardless of what insurance you have, or if you have no insurance at all, **Verastem Cares™** has programs to support you.

Verastem Cares™ will work with your healthcare provider to help you get access to treatment.

HAVE A QUESTION? WANT TO LEARN MORE?

Just call us. Our dedicated team is here to help.

Phone: 866-351-8372 8:00 am to 8:00 pm ET Monday-Friday

Click to read full [Prescribing Information](#) and [Patient Information](#).



© 2025 Verastem, Inc. All Rights Reserved.
AF-US-2025-0009 05/25

AVMAPKI and FAKZYNJA are trademarks of Verastem, Inc.

Verastem Cares is a trademark of Verastem, Inc.

For US patients and caregivers 18 years or older

 **AVMAPKI™ FAKZYNJA™ CO-PACK**
(avutemetinib capsules; defactinib tablets) 0.8 mg/200 mg