

REIMAGINE

RECURRENT LGSOC



Treatment Tracker:

A tool for tracking your side effects during treatment and talking to your healthcare provider

THE FIRST FDA-APPROVED TREATMENT SPECIFICALLY for women who have low-grade serous ovarian cancer (LGSOC) that has come back (recurrent), and LGSOC with an abnormal *KRAS* gene, and previously been treated with medicine for their cancer.

It is not known if Avmapki Fakzynja Co-Pack is safe and effective in children.

IMPORTANT SAFETY INFORMATION

Avmapki Fakzynja Co-Pack may cause serious side effects, including eye problems, severe skin reactions, liver problems, and muscle problems such as rhabdomyolysis. These are not all the possible side effects of Avmapki Fakzynja Co-Pack. Call your doctor for medical advice about side effects.

Please see additional Important Safety Information throughout this brochure and discuss with your doctor.

Click here for full [Prescribing Information](#) and [Patient Information](#).

 **AVMAPKI™ FAKZYNJA™ CO-PACK**
(avutometinib capsules; defactinib tablets) 0.8 mg; 200 mg

This guide is designed to help you track your side effects during treatment and talk to your healthcare provider about your treatment. Just add your questions and notes, and bring it to your next appointment.

What is Avmapki Fakzynja Co-Pack?

Avmapki Fakzynja Co-Pack is a prescription medicine used to treat adults who have:

- low-grade serous ovarian cancer (LGSOC) that has come back (recurrent), **and**
- LGSOC with an abnormal *KRAS* gene, **and**
- previously been treated with medicine for their cancer.

It is not known if Avmapki Fakzynja Co-Pack is safe and effective in children.



Let's get started: Your medical history and plans for pregnancy

Your doctor needs to know about any health problems you have, especially if they affect your eyes, skin, or liver. It's also important to let your healthcare provider know if you are pregnant, breastfeeding, or planning to become pregnant.

What are all your medical conditions?

Please list questions and concerns about pregnancy:

IMPORTANT SAFETY INFORMATION (CONT'D)

Before taking Avmapki Fakzynja Co-Pack, tell your healthcare provider about all of your medical conditions, including if you:

- have a history of eye or vision problems
- have had severe skin reactions
- have liver problems
- are pregnant or plan to become pregnant. Avmapki Fakzynja Co-Pack can harm your unborn baby.
 - **Females who are able to become pregnant:**
 - Your healthcare provider should do a pregnancy test before you start treatment with Avmapki Fakzynja Co-Pack.
 - You should use effective birth control (contraception) during treatment and for 1 month after the last dose of Avmapki Fakzynja Co-Pack.
 - Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with Avmapki Fakzynja Co-Pack.
 - **Males with a female partner who is able to become pregnant:**
 - You should use effective birth control (contraception) during treatment and for 4 months after the last dose of Avmapki Fakzynja Co-Pack.
- are breastfeeding or plan to breastfeed. It is not known if Avmapki Fakzynja Co-Pack passes into your breast milk. Do not breastfeed during treatment and for 2 weeks after the last dose of Avmapki Fakzynja Co-Pack.

Please see additional Important Safety Information throughout this brochure and discuss with your doctor.



Let's get started: About you

Avmapki Fakzynja Co-Pack may affect the way other medicines work, and other medicines may affect how Avmapki Fakzynja Co-Pack works. It's important to tell your healthcare provider all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Especially tell your healthcare provider if you take:

Proton pump inhibitors (PPIs), such as omeprazole, pantoprazole, etc.

H2 receptor antagonists (H2 blockers), such as cimetidine, ranitidine, etc.

A blood thinner called warfarin

Please check the applicable boxes above, and additionally list all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements:

IMPORTANT SAFETY INFORMATION (CONT'D)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Avmapki Fakzynja Co-Pack may affect the way other medicines work, and other medicines may affect how Avmapki Fakzynja Co-Pack works.

Especially tell your healthcare provider if you take:

- proton pump inhibitors (PPIs) and H2 receptor antagonists
- a blood thinner called warfarin

Know the medicines you take. Keep a list of them with you to show your healthcare provider and pharmacist when you get a new medicine.



Next up: Taking Avmapki Fakzynja Co-Pack

How should I take Avmapki Fakzynja Co-Pack?

- Take Avmapki Fakzynja Co-Pack exactly as directed by your healthcare provider
- Do not change your dose or stop taking Avmapki Fakzynja Co-Pack without talking to your healthcare provider
- If you develop certain side effects, your healthcare provider may change your dose, temporarily stop, or completely stop your treatment with Avmapki Fakzynja Co-Pack
- Take Avmapki capsules and Fakzynja tablets with food
- If you vomit after taking Avmapki capsules or Fakzynja tablets, do not take an additional dose. Take the next scheduled dose as prescribed by your healthcare provider



Avmapki capsules:

- Take each dose at the same time
- Swallow the capsules whole. Do not chew, break, or open the capsules
- If you miss a dose of Avmapki by more than 24 hours, skip the missed dose and take the next scheduled dose as prescribed by your healthcare provider. Do not take two doses at the same time to make up for a missed dose



Fakzynja tablets:

- Swallow the tablets whole. Do not chew, break, or crush the tablets
- If you take antacids, take Fakzynja tablets 2 hours before or 2 hours after taking the antacid
- If you miss a dose of Fakzynja by more than 6 hours, skip the missed dose and take the next scheduled dose as prescribed by your healthcare provider. Do not take two doses at the same time to make up for a missed dose

Product images shown are not actual size.

IMPORTANT SAFETY INFORMATION (CONT'D)

Before and during treatment with Avmapki Fakzynja Co-Pack, your healthcare provider will give you medicines to apply to your face, scalp, neck, upper chest and upper back, and antibiotic medicines to take by mouth to help prevent skin reactions.

What should I avoid while taking Avmapki Fakzynja Co-Pack?

- Limit your time in the sun. Avmapki Fakzynja Co-Pack can make your skin sensitive to the sun (photosensitivity) and the light from sunlamps and tanning beds. You may burn more easily and get a severe sunburn. Apply daily sunscreen with a Sun Protection Factor (SPF) of 30 or greater and wear a hat and clothes that cover your skin to protect against sunburn. Talk to your healthcare provider if you get a sunburn.

Please see additional Important Safety Information throughout this brochure and discuss with your doctor.



Next up: Monitoring your eye health

While taking Avmapki Fakzynja Co-Pack, your healthcare provider will request eye exams to monitor your eyes and vision.

Please check the boxes of the symptoms you are experiencing and review them with your healthcare provider at each visit. Please notify your healthcare provider immediately of any new or worsening symptoms.

Changes in vision, such as blurred vision, double vision, or vision loss	Eye pain
Dry eye(s)	New or increased floaters (small dark spots or squiggly lines that float across your vision)
Eye and eyelid inflammation, including redness, swelling, and itching	Seeing flashes of light
Light hurting your eyes	


Please add any additional eye-related symptoms, comments, or questions here:

IMPORTANT SAFETY INFORMATION (CONT'D)

What are the possible side effects of Avmapki Fakzynja Co-Pack?

Avmapki Fakzynja Co-Pack may cause serious side effects, including:

- **eye problems.** Eye problems are common with Avmapki Fakzynja Co-Pack and can be severe. Your healthcare provider will send you to see an eye care professional before starting treatment, during treatment, and for any new or worsening eye problems. Tell your healthcare provider if you get any new or worsening eye problems including:
 - changes in vision, such as blurred vision, double vision, or vision loss
 - dry eye(s)
 - eye and eyelid inflammation, including redness, swelling, and itching
 - eye pain
 - new or increased floaters (small dark spots or squiggly lines that float across your vision)
 - light hurting eyes or seeing flashes of light



Next up: Monitoring your skin health

Before and during treatment with Avmapki Fakzynja Co-Pack, your healthcare provider will give you medicines to apply to your face, scalp, neck, upper chest and upper back, and antibiotic medicines to take by mouth to help prevent skin reactions.

Please check the boxes of the symptoms you are experiencing and review them with your healthcare provider at each visit. Please notify your healthcare provider immediately of any new or worsening symptoms.

Skin rash or acne	Redness or swelling of your face, hands, or soles of your feet
Dry skin	Blisters or sores in your mouth
Itchy skin	Peeling of your skin
Blisters on your skin	Skin infection

Please add any additional skin-related symptoms, comments, or questions here:

IMPORTANT SAFETY INFORMATION (CONT'D)

What are the possible side effects of Avmapki Fakzynja Co-Pack?

Avmapki Fakzynja Co-Pack may cause serious side effects, including:

- **severe skin reactions.** Skin reactions are common with Avmapki Fakzynja Co-Pack and can be severe or life threatening. Tell your healthcare provider if you develop any new or worsening skin reactions including:
 - skin rash or acne
 - dry skin
 - itching
 - blisters on your skin
 - redness or swelling of your face, hands, or soles of your feet
 - blisters or sores in your mouth
 - peeling of your skin
 - skin infection

Please see additional Important Safety Information throughout this brochure and discuss with your doctor.



Next up: Monitoring your liver health

While taking Avmapki Fakzynja Co-Pack, your healthcare provider will request blood tests to monitor your liver health.

Please check the boxes of the symptoms you are experiencing and review them with your healthcare provider at each visit. Please notify your healthcare provider immediately of any new or worsening symptoms.

Yellowing of your skin or the white part of your eyes	Nausea or vomiting
Itchy skin	Upper right side stomach pain
Feeling very tired	Dark or tea colored urine
Flu-like symptoms	Bleeding or bruising more easily than normal

Please add any additional symptoms, comments, or questions here:

IMPORTANT SAFETY INFORMATION (CONT'D)

What are the possible side effects of Avmapki Fakzynja Co-Pack?

Avmapki Fakzynja Co-Pack may cause serious side effects, including:

- **liver problems.** Liver problems are common with Avmapki Fakzynja Co-Pack and can be severe. Treatment with Avmapki Fakzynja Co-Pack can increase the level of aspartate aminotransferase, alanine aminotransferase, and bilirubin in your blood. Your healthcare provider will do blood tests before starting each treatment cycle and during treatment to check your liver function. Tell your healthcare provider if you get any signs and symptoms of severe liver problems including:
 - yellowing of your skin or the white part of your eyes
 - itchy skin
 - feeling very tired
 - flu-like symptoms
 - nausea or vomiting
 - upper right side stomach pain
 - dark or tea colored urine
 - bleeding or bruising more easily than normal.

Please see additional Important Safety Information throughout this brochure and discuss with your doctor.



Next up: Monitoring other aspects of your physical health

While taking Avmapki Fakzynja Co-Pack, your healthcare provider will monitor your muscle health.

Please check the boxes of the symptoms you are experiencing and review them with your healthcare provider at each visit. Please notify your healthcare provider immediately of any new or worsening symptoms.

Weakness or difficulty moving arms and legs

Muscle or bone aches or pain that does not go away

Dark, red colored urine

Decreased urine output

Please add any additional symptoms, comments, or questions here:

IMPORTANT SAFETY INFORMATION (CONT'D)

What are the possible side effects of Avmapki Fakzynja Co-Pack?

Avmapki Fakzynja Co-Pack may cause serious side effects, including:

- **muscle problems (rhabdomyolysis).** Avmapki Fakzynja Co-Pack may cause muscle problems that can be severe. Treatment with Avmapki Fakzynja Co-Pack can increase the level of an enzyme in your blood called creatine phosphokinase (CPK) and can be a sign of muscle damage. Your healthcare provider will perform blood tests to check your levels of CPK before starting each treatment cycle and during treatment. Tell your healthcare provider if you get any signs or symptoms of increased CPK including:
 - weakness or difficulty moving arms and legs
 - muscle or bone aches or pain that does not go away
 - dark, red colored urine or decreased urine output.

Please see additional Important Safety Information throughout this brochure and discuss with your doctor.



Next up: Monitoring your overall health

It's important that you tell your healthcare provider about any problems or discomfort that you may be experiencing during treatment.

Please check the boxes of the symptoms you are experiencing and review them with your healthcare provider at each visit. Please notify your healthcare provider immediately of any new or worsening symptoms.

Nausea	Heartburn (acid reflux)
Tiredness	Constipation
Diarrhea	Shortness of breath
Swelling in the body (edema)	Cough
Vomiting	Urinary tract infection
Stomach-area pain	Abnormal blood tests

These are not all of the possible side effects of Avmapki Fakzynja Co-Pack. For more details, please see the Important Safety Information throughout this treatment tracker and the full Prescribing Information.

Please add any additional symptoms, comments, or questions here:

IMPORTANT SAFETY INFORMATION (CONT'D)

The most common side effects of Avmapki Fakzynja Co-Pack include:

- nausea
- tiredness
- increased aspartate aminotransferase
- diarrhea
- swelling in the body (edema)
- decreased hemoglobin
- increased alanine aminotransferase
- vomiting
- increased blood bilirubin
- increased triglycerides
- stomach-area pain
- acid reflux
- increased alkaline phosphatase
- decreased platelets
- constipation
- shortness of breath
- cough
- urinary tract infection
- decreased lymphocytes
- decreased neutrophils
- proteinuria

Please see additional Important Safety Information throughout this brochure and discuss with your doctor.

